ESRF	Experiment title: The evolutionary origin of metamorphosis in tetrapods: clues from palatal osteohistology in temnospondyls	Experiment number: LS-3109
Beamline:	Date of experiment:	Date of report:
BM18	from: 26.10.2022 to: 29.10.2022	9.3.2023
Shifts:	Local contact(s):	Received at ESRF:
9	Kathleen Dollman, Vincent Fernandez	10.3.2023

Names and affiliations of applicants (* indicates experimentalists):

Pérez, Celeste Marina*. Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Berlin, Germany

Anne-Claire Fabre*. Naturhistorisches Museum Bern, Bern, Switzerland; Institute of Ecology and Evolution, University of Bern, Bern, Switzerland; Life Sciences Department, Vertebrates Division, Natural History Museum, London, United Kingdom

Donald Davesne*. Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Berlin, Germany

Nadia Fröbisch. Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Berlin, Germany; Humboldt-Universität zu Berlin, Berlin, Germany

Report:

During the nine shifts, we could scan 19 salamander specimens and four fossil temnospondyls. Scans at $4.2 \mu m$ and $0.85 \mu m$ were made.

The scanning of the salamanders took us longer than expected because we had problems finding the regions of interest (ROIs) for the scans at 0.85µm, which requiered us to perform numerous test scans. Because of this delay, we were not able to scan all the fossils that are essential for the project. Therefore, one of us (C.P.) returned to the ESRF in December 2022 to successfully complete the remaining scans in three shifts (experiment IH-LS-3441).

To date, the processed scans are still unavailable, so we have not been able to properly evaluate whether the data collected at the ESRF will allow us to answer the research questions detailed in the proposal. However, preliminary reconstructions made during the scanning at the ESRF look promising; at 0.85µm we could see histological features in the extant specimens and confirm that the preservation of most fossils was good at the histological level.